

BROWN-GITLER SPECTRA AT $BP\langle 2 \rangle$

John KLIPPENSTEIN

*Department of Mathematics, Middlesex College, The University of Western Ontario, London, Canada
N6A 5B7*

Received 30 December 1986

Revised 14 April 1987

Brown-Gitler spectra for the homology theory associated to $BP\langle 2 \rangle$ are constructed. Complexes adapted to the new Brown-Gitler spectra are constructed and a spectral sequence converging to stable maps into these spectra is examined.

AMS (MOS) Subj. Class.: Primary 54P42, 55N35

Brown-Gitler spectra $BP\langle 2 \rangle$

In [1], Brown and Gitler constructed certain spectra $B(k)$ with the property that $H^*B(k) \cong A/I_k$ for certain ideals I_k over the Steenrod algebra A . In [6], Goerss, Jones and Mahowald constructed what they called generalized Brown-Gitler spectra at $BP\langle 1 \rangle$, written $BP\langle 1 \rangle^k$, with the property that $H^*BP\langle 1 \rangle^k \cong A/I_k + J_1$ where J_1 is the ideal with the property that $H^*BP\langle 1 \rangle \cong A/J_1$. This paper is devoted to constructing Brown-Gitler spectra at $BP\langle 2 \rangle$ with $H^*BP\langle 2 \rangle^k \cong A/I_k + J_2$ where $H^*BP\langle 2 \rangle \cong A/J_2$.

Although the original Brown-Gitler spectra have been put to a variety of uses in homotopy theory, the applications of generalized Brown-Gitler spectra have been restricted to splittings of $bo \wedge bo$ (see [9]), $BP\langle 1 \rangle \wedge BP\langle 1 \rangle$ (see [7]), and $BP\langle 1 \rangle \wedge BP\langle n \rangle$ (see [8]). In fact if J_n is an ideal satisfying $H^*BP\langle n \rangle \cong A/J_n$ (one should interpret $BP\langle -1 \rangle$ as $H\mathbb{Z}/p$, $BP\langle 0 \rangle$ as $H\mathbb{Z}_p^\wedge$, and J_{-1} as the empty set), then $H^*(BP\langle n \rangle \wedge BP\langle n \rangle) \cong \bigoplus_{k \geq 0} \Sigma^{2(p-1)k} H^*BP\langle n \rangle \otimes A/I_k + J_n$ and one hopes to be able to realize $A/I_k + J_n$ as the mod p cohomology of a spectrum $BP\langle n \rangle^k$ and then realize this as a splitting of spectra. This is the motivation for constructing $BP\langle 2 \rangle^k$.

Having fixed k , the first task is to construct an acyclic resolution of $A/I_k + J_n$. $A/I_k + J_n \leftarrow D_0^n \rightarrow D_1^n \leftarrow D_2^n \leftarrow \dots$. Brown and Gitler [1] used the lambda algebra to produce the corresponding resolution; Goerss, Jones and Mahowald [6] used a modification of the lambda algebra. The appropriate modification for $BP\langle 2 \rangle$ is apparent in [6] and little will be said here about the details needed to get the acyclic resolution,

$$A/I_k + J_2 \leftarrow D_0^2 \leftarrow D_1^2 \leftarrow D_2^2 \leftarrow \dots$$

Along with this, one wants to have the resolution realized by maps of spectra,

$$K_0^n \xrightarrow{d_0} \Sigma K_1^n \xrightarrow{d_1} \Sigma^2 K_2^n \rightarrow \cdots$$

Since in [1] the resolution was free over the Steenrod algebra, they got the maps of spectra for free. This was not the case in either [6] or this paper. In our resolution the first three terms are direct sums of suspensions of $H^*BP\langle 2 \rangle$, $H^*BP\langle 1 \rangle$ and $H^*H\mathbb{Z}_p^\wedge$ respectively. The only new constructions needed are maps $S^j: BP\langle 2 \rangle \rightarrow \Sigma^{2(p-1)}BP\langle 1 \rangle$ with the property that in cohomology $S^{j*}(1) = \chi(P^j)$, where 1 and $\chi(P^j)$ represent their respective equivalence classes in A/J_1 and A/J_2 . This is a consequence of the fact that the Adams spectral sequence for $[BP\langle 2 \rangle, BP\langle 1 \rangle]$ collapses to its E_2 term, as proved in [8].

Conceptually, the next step would be to construct a tower of spectra with the following properties.

Theorem A. *For any n between -1 and 2 there is a tower of spectra*

$$X_{q+1} \xrightarrow{p_1} X_q \rightarrow \cdots \rightarrow X_1 \xrightarrow{p_0} X_0$$

with the property that

- (a) *there are maps $e_1: X_q \rightarrow \Sigma K_{q+1}^n$ where K_{q+1}^n are the spectra mentioned in the previous paragraph,*
- (b) $X_0 = K_0^n = BP\langle n \rangle$, $e_0 = d_0$,
- (c) $K_{q+1}^n \xrightarrow{i_{q+1}} X_{q+1} \xrightarrow{p_q} X_q \xrightarrow{e_q} \Sigma K_{q+1}^n$,
is a cofiber sequence (defining i_{q+1}),
- (d) $e_q i_q = d_q$, $i_0 = Id$,
- (e) *For any CW complex Z , the induced map of homology theories $e_q^*: (X_q)_m Z \rightarrow (K_{q+1}^n)_m Z$ is zero for $m \leq 2p(k+1) - 1$ and $q \geq n$.*

This was proved in [6] for $-1 \leq n \leq 1$ (one should identify $BP\langle -1 \rangle$ with $H\mathbb{Z}/p$ and $BP\langle 0 \rangle$ with $H\mathbb{Z}_p^\wedge$). Given Theorem A, the $BP\langle 2 \rangle^k$ are defined to be the homotopy inverse limit of the X_q tower in the theorem when $n = 2$.

Theorem B. *For each $k \geq 0$ and prime p , there is a p -complete spectrum $BP\langle 2 \rangle^k$ and a map $w: BP\langle 2 \rangle^k \rightarrow BP\langle 2 \rangle$ such that*

- (a) $H^*BP\langle 2 \rangle^k \cong A/A\{\beta, \beta P^1 \beta P^{p+1}, \chi P^j \mid j > k\} \cong A/I_k + J_2$
- (b) $\{\text{Im } w_*: BP\langle 2 \rangle_m^k Z \rightarrow BP\langle 2 \rangle_m Z\} \subseteq \bigcap_{j > k} \ker S_*^j$.

*If Z is a CW complex with $H\mathbb{Z}_p^\wedge * Z$ an \mathbb{F}_p vector space and $m \leq 2p(k+1) - 1$ then $\text{Im } w_*$ is all of $\bigcap_{j > k} \ker S_*^j$.*

Partial Proof. To prove part (a), apply H^* to the tower of fibrations in Theorem A. The spectral sequence associated to the unravelled exact couple converges to

$H^*\text{holim}_q X_q = H^*BP\langle 2 \rangle^k$ and has $E_{s,*}^1 \cong H^*K_s^2 \cong D_s^2$. Since $0 \leftarrow A/I_k + J_2 \leftarrow D_0^2 \leftarrow D_1^2 \leftarrow \dots$ was an acyclic resolution, the E^2 term of this spectral sequence has $E_0^2 = \text{Coker } H^*d_0$ and $E_i^2 = 0$ for $i \geq 1$. Thus $H^*BP\langle 2 \rangle^k$ is as claimed in (a).

The map $w: BP\langle 2 \rangle^k \rightarrow BP\langle 2 \rangle$ is just

$$BP\langle 2 \rangle^k = \text{holim}_q X_q \rightarrow X_0 = BP\langle 2 \rangle$$

which can be factored as

$$BP\langle 2 \rangle^k \rightarrow X_1 \xrightarrow{p_0} BP\langle 2 \rangle.$$

Since

$$X_1 \rightarrow BP\langle 2 \rangle \xrightarrow{\bigcap_{j>k} S^j} \bigcup_{j>k} \Sigma^{2(p-1)j} BP\langle 1 \rangle = \Sigma K_1^2$$

is a cofibration we get the first part of assertion (b). The proof of the last part of (b) requires some knowledge of how Theorem A is proved and will appear in Section 3.

Finally taking the tower of fibrations created in Theorem A, smashing with any spectrum Y and taking stable homotopy gives a spectral sequence converging to $BP\langle n \rangle_*^k Y$ with

$$E_{s,t}^1(Y \wedge BP\langle n \rangle) \cong (K_s^n)_{t-s} Y = \pi_{t-s}(K_s \wedge Y).$$

When $n=0$ or -1 and Y is the suspension spectrum of a space then part (e) of Theorem A implies that

$$E_{s,t}^1(Y \wedge BP\langle n \rangle) \cong E_{s,t}^\infty(Y \wedge BP\langle n \rangle) \quad \text{for } t-s \leq 2p(k+1)-2.$$

When $n=1$ this becomes $E_{s,t}^2(Y \wedge BP\langle n \rangle) \cong E_{s,t}^\infty(Y \wedge BP\langle n \rangle)$ for $t-s \leq 2p(k+1)-2$. For Brown-Gitler spectra at $BP\langle 2 \rangle$ we get

Corollary C. *If Y is the suspension spectrum of a space then in the spectral sequence converging to $BP\langle 2 \rangle_*^k Y$,*

$$E_{s,t}^1(Y \wedge BP\langle 2 \rangle) \cong E_{s,t}^\infty(Y \wedge BP\langle 2 \rangle) \quad \text{for all } t-s \leq 2p(k+1)-2.$$

The dual to this situation involves applying the functor $[Y,]$ to the tower of Theorem A to get a spectral sequence converging to $[Y, BP\langle n \rangle^k]^{s-t}$ with

$$E_{t,s}^{s,t}(Y, BP\langle n \rangle) \cong [Y, K_s^n]^{s-t}.$$

In order to get the collapsing result analogous to Corollary C, [6] introduced the notion of spacelike spectra.

Definition. Let Y be a finite CW spectrum. Y is said to be *spacelike of dimension n* if there is a spectrum T and a map $f: T \rightarrow Y$ so that

- (i) $T = \Sigma^n DZ$ where Z is a finite CW complex, and
- (ii) $f^*: H^*Y \rightarrow H^*T$ is injective.

Corollary D. *If Y is spacelike of dimension n , then in the spectral sequence converging to $[Y, BP\langle 2 \rangle^k]^{s-t}$, for all $s - t \geq n - 2p(k + 1) + 2$*

$$E_3^{s,t} \cong E_\infty^{s,t} \quad \text{and} \quad E_1^{s,t} \cong E_\infty^{s,t} \quad \text{for } s \geq 3.$$

Although conceptually the construction of the X_q tower comes after realizing the D_n^2 resolution, performing the construction requires the use of some complicated machinery and while constructing the tower one must simultaneously construct maps from the spectra needed for the $BP\langle 2 \rangle^k$ construction to those that were used in the construction of the original Brown-Gitler spectra.

In Section 1 the acyclic resolution of $M(2, k)$ is given and realized. Section 2 contains the definition and construction of adapted complexes, the complicated machinery referred to in the previous paragraph. Section 3 gives the proof of Theorem A and Theorem B while Section 4 concludes with a proof of Corollary D.

Throughout this paper we are working with a fixed prime p and a fixed positive integer k . Homology is always with mod p coefficients unless otherwise specified.

In this paper, references to earlier results about Brown-Gitler spectra and their construction are almost always to [6]. This is for convenience since that is where the results are stated in the form we need. In fact all results about Brown-Gitler spectra at $BP\langle -1 \rangle = H\mathbb{Z}/p$ are due to [1] when $p = 2$ and [3] when p is odd. Brown-Gitler spectra at $BP\langle 0 \rangle = H\mathbb{Z}_p^\wedge$ are due to Mahowald [9] when $p = 2$ and Kane [7] for p odd and also to Goerss [4] and Shimamoto [10].

1. Acyclic resolutions

In this section we collect all the definitions and results from [6] regarding the construction and realization of the acyclic resolutions that we need. Their work made extensive use of the A algebra which we will not do, since their work is sufficient to construct the acyclic resolution we need and the A algebra's work is done once they have been constructed. Throughout this section, if the prime you have chosen is $p = 2$, replace P^i by Sq^{2^i} and β by Sq^1 .

For $n \geq 0$, let E_n be the Hopf subalgebra of the Steenrod algebra A , generated by the Milnor elements Q_0, Q_1, \dots, Q_n , let \bar{E}_n be the augmentation ideal and J_n the left A ideal generated by \bar{E}_n . Notice that $H^*BP\langle n \rangle \cong A/J_n$. For $n \geq 1$, let $M_n(k) = A/I_k + J_n$ (where $J_{-1} = \emptyset$). Theorem 2.6 of [6] gives an acyclic resolution of $M_n(k)$ by A modules

$$\cdots \rightarrow D_{q+1}^n \xrightarrow{d_q^*} D_q^n \rightarrow \cdots \rightarrow D_1^n \xrightarrow{d_0^*} D_0^n \xrightarrow{\epsilon} M_n(k) \rightarrow 0$$

for $n = 0$ and 1. Their proof works equally well for $n = 2$. One should notice that

$D_0^2 = H^*BP\langle 2 \rangle$, $D_1^2 = \bigvee_{j>k} \Sigma^{qj} H^*BP\langle 1 \rangle$ and D_2^2 is a sum of $H^*BP\langle 0 \rangle$'s while for $n \geq 3$, D_n^2 is a free module and a direct summand of D_{n-1}^2 which is also free. In general, D_m^n is a direct sum of copies of $H^*BP\langle \max(-1, n-m) \rangle$ and the set indexing the copies is a subset of the set indexing copies of D_m^{n-r} if $r > 0$. Thus there are quotient maps $\theta_q^*: D_q^{-1} \rightarrow D_q^n$ so that

$$\begin{array}{ccc} D_{q+1}^{-1} & \xrightarrow{d_q^*} & D_q^{-1} \\ \downarrow \theta_{q+1}^* & & \downarrow \theta_q^* \\ D_{q+1}^n & \longrightarrow & D_q^n \end{array}$$

commutes and for $q \geq 2$, θ_q^* is a projection onto a direct summand. We can choose spectra K_q^n such that $H^*K_q^n \cong D_q^n$ and get the following:

Proposition 1.1. *For $q \geq 0$, $0 \leq n \leq 2$, there exist maps $\theta_q: K_q^n \rightarrow K_q^{-1}$ and $d_q: K_q^n \rightarrow \Sigma K_{q+1}^n$ so that*

- (i) $H^*(\theta_q): D_q^{-1} \rightarrow D_q^n$ is just the θ_q^* mentioned above,
- (ii) $H^*(d_q): D_{q+1}^n \rightarrow D_q^n$ is the differential d_q^* in the acyclic resolution,

$$\text{(iii)} \quad \begin{array}{ccc} K_q^n & \xrightarrow{d_q} & \Sigma K_{q+1}^n \\ \downarrow \theta_q & & \downarrow \theta_{q+1} \\ K_q^{-1} & \xrightarrow{d_q} & \Sigma K_{q+1}^{-1} \end{array} \quad \text{commutes,}$$

- (iv) if $q \geq n+1$ there is a map $s_q: K_q^{-1} \rightarrow K_q^n$ so that $s_q \theta_q$ is the identity.

Proof. Cases $n=0$ and 1 were done in Proposition 2.7 of [6]. For $n=2$, parts (i) and (iv) follow immediately from the structure of the D_q^n 's mentioned earlier in this paragraph. Since for $q \geq 2$, K_q^2 is a wedge of Eilenberg-MacLane spectra, the only thing that needs to be done is the construction of d_0 and d_1 . To get d_0 , we want to realize d_0^* . Looking at d_0^* 's effect on just one factor (see [6]), we have

$$\begin{array}{ccc} \Sigma^{2(p-1)j} H^*BP\langle 1 \rangle & \rightarrow \bigoplus_{i>k} \Sigma^{2(p-1)i} H^*BP\langle 1 \rangle & \xrightarrow{d_0^*} H^*BP\langle 2 \rangle \\ a \longmapsto & & a \cdot \chi(P^j). \end{array}$$

This is well defined since the formula

$$P^j Q_i - Q_i P^j = Q_{i+1} P^{j-p^i}$$

implies that $P^j Q_0$ and $P^j Q_1$ are in $\bar{E}_2 \cdot A$ so that $Q_0 \chi P^j$ and $Q_1 \chi P^j$ are zero in $H^*BP\langle 2 \rangle \cong A/A \cdot \bar{E}_2$. Since, as proved in [8], the Adams spectral sequence for

$[BP\langle 2 \rangle, BP\langle 1 \rangle]$ collapses to its E_2 term, this element of $E_2^{0,2(p-1)j} = \text{Hom}^{2(p-1)j}(H^*BP\langle 1 \rangle, H^*BP\langle 2 \rangle)$ gives a map $S^j = BP\langle 2 \rangle \rightarrow \Sigma^{2(p-1)j} BP\langle 1 \rangle$ making

$$\begin{array}{ccc} BP\langle 2 \rangle & \xrightarrow{S^j} & \Sigma^{2(p-1)j} BP\langle 1 \rangle \\ \downarrow \rho & & \downarrow \rho \\ H\mathbb{Z}/p & \xrightarrow{\chi^{P^j}} & \Sigma^{2(p-1)j} H\mathbb{Z}/p \end{array}$$

commute. The map d_1 is handled in the same way. \square

2. Adapted complexes

Let B a subalgebra of A and E a ring spectrum with $H^*E = A//B = A/\bar{A}\bar{B}$. Write $N(k)$ for $A/A\{\bar{B}, w_k\}$ where $w_k = \{\chi^{P^i} \mid i > k\}$ and let $1: E \rightarrow H$ be the generator of H^*E over A . If Z is a finite CW complex and $h' \in E_m Z$ with $m \leq 2p(k+1)-1$ and $h: \Sigma^m DZ \rightarrow E$ is the dual of h' , then we say that (Z, h') is *adapted to $N(k)$* if

$$A\{\bar{B}, w_k\} \rightarrow A \xrightarrow{h^* \cdot 1^*} H^* \Sigma^m DZ$$

is exact. The degree of the adapted complex is the degree m of h' in $E_* Z$. The aim of this section is to prove the following theorem.

Theorem 2.1. *For each $k \geq 0$, there is a finite CW complex Z_k and $h'_k \in BP\langle 2 \rangle_m Z_k$ with $m = 2pk + 3$ so that (Z_k, h'_k) is adapted to $M_2(k)$. Also h'_k can be chosen so that $d_{0*} h'_k = 0$ where d_0 is as in Proposition 1.1.*

Let $R = \{\chi^{P^I} \mid Q_2 Q_1 Q_0 P^I \neq 0, I \text{ admissible}, i_1 \geq k\}$. If $\varepsilon: A \rightarrow M_2(k)$ is the projection map, then $\varepsilon(R)$ forms a vector space basis for $M_2(k)$. Lemma 3.6 of [6] reduces the construction of an adapted complex for $M_2(k)$ to proving the next proposition.

Proposition 2.2. *For each $P \in R$ there is a CW complex Z_P and $\bar{h}_P \in H\mathbb{Z}/p_{2pk+3} Z_P$ with*

- (1) $\chi(P)^* \bar{h}_P \neq 0$,
- (2) $\bar{h}_P = 1_* h'_P$ for some $h'_P \in BP\langle 2 \rangle_* Z_P$ where 1_* is the reduction

$$BP\langle 2 \rangle_* Z_P \xrightarrow{1_*} H\mathbb{Z}/p_* Z_P.$$

This will suffice since their lemma tells us that $Z = \bigvee_{P \in R} Z_P$ and $h' = \bigoplus_{P \in R} h'_P$ will be adapted to $M_2(k)$.

Proof of Proposition 2.2. Take any $\chi^{P^I} \in R$, so $Q_2 Q_1 Q_0 P^I \neq 0$, $I = (i_1, \varepsilon_1, i_2, \varepsilon_2, \dots, i_m, \varepsilon_m)$ is admissible and $i_1 \geq k$. $Q_2 Q_1 Q_0 P^I \neq 0$, implies $\varepsilon_1 = 0 = \varepsilon_2$. Set

$$e = 2i_1 - \sum_{j=2}^m 2(p-1)i_j - \sum_{j=3}^m \varepsilon_j, \quad t = \sum_{i=3}^m \varepsilon_i \quad \text{and} \quad s = \frac{1}{2}(e-t).$$

The Z associated to χP^I will be a subcomplex of

$$C(s, t+3) = \bigwedge_s \mathbb{C}P^\infty \wedge \bigwedge_{t+3} B\mathbb{Z}/p,$$

the s -fold smash product of $\mathbb{C}P^\infty$ smashed with the $(t+3)$ -fold smash product of $B\mathbb{Z}/p$. Then $H^{2s+t+3}C(s, t+3)$ is the first nonzero group and is generated by a single nonzero element c .

Let pr be the vector space map

$$\text{pr}: H^*C(s, t+3) \rightarrow (H^*C(s, t+3)/\bar{E}_2 \cdot H^*C(s, t+3)).$$

Now s and t have been chosen so that $Q_2Q_1Q_0P^I C \neq 0$. This implies that $P^I c$ is not in $\bar{E}_2 \cdot H^*C(s, t+3)$ so that $v = \text{pr}(P^I c) \neq 0$. Dualizing we get

$$(P^I)^* \text{pr}^*(v^*) \neq 0 \quad \text{in } H_*C(s, t+3).$$

Let X_I be a finite skeleton of $C(s, t+3)$ containing the $2p(k+1) + |Q_0| + |Q_1| + |Q_2|$ skeleton and such that $H\mathbb{Z}_p^{\wedge} X_I$ is an F_p vector space. Then $H\mathbb{Z}_p^{\wedge} X_I \rightarrow H\mathbb{Z}/p_* X_I$ is a monomorphism. Let $j = 2pk - 2pi_1$ and $Z_I = \Sigma^j X_I$. Let $c \in H^{j+2s+t+3} Z_I$ be the generator. Then we still have $\text{pr}(P^I c) \neq 0$ in $H^*Z_I/\bar{E}_2 \cdot H^*Z_I$. Setting $\bar{h}_I = \text{pr}^*(\text{pr}(P^I c))^* \in H_{2pk+3} Z_I$ we find that $(P^I)^* \bar{h}_I \neq 0$. All that we have left to show is that \bar{h}_I is in the image of the reduction map $1_*: \text{BP}\langle 2 \rangle_* Z_I \rightarrow H_* Z_I$. This follows from putting together some facts from [6] and [5]. In [5], Goerss proves that the Adams spectral sequence converging to $\text{BP}\langle 2 \rangle_* C(s, t)$ collapses to its E_2 term. This ensures that in the required range of dimensions the Adams spectral sequence converging to $\text{BP}\langle 2 \rangle_* Z_I$ collapses to its E_2 term. As pointed out in [6], one easily checks that this E_2 term satisfies

$$\begin{aligned} \text{Ext}_A^0(H^*\text{BP}\langle 2 \rangle \otimes H^*Z_I, \mathbb{Z}/p) &\cong \text{Hom}_{E_2}^*(H^*Z_I, \mathbb{Z}/p) \\ &\cong (H^*Z_I/\bar{E}_2 \cdot H^*Z_I)^* \end{aligned}$$

and the composite.

$$\text{BP}\langle 2 \rangle_* Z_I \rightarrow E_\infty^0 \cong (H^0 Z_I/\bar{E}_2 \cdot H^*Z_I)^* \rightarrow H\mathbb{Z}/p_* X$$

is the reduction induced by $1: \text{BP}\langle 2 \rangle \rightarrow H\mathbb{Z}/p$. From this we can conclude that $\bar{h}_I = \text{pr}^*(\text{pr}(P^I c))^*$ is in the image of this reduction map; so choose some h'_I with $1_* h'_I = \bar{h}_I$.

Having constructed the adapted complex mentioned in Theorem 2.1, all that remains to be proved of that theorem is that $d_{0*} h'_I = 0$. The fact that $P^I c$ generated a free E_2 module in H^*Z_I means that h'_I will generate a submodule of $\text{BP}\langle 2 \rangle_* Z_I$ with trivial p , v_1 and v_2 actions. This means that if we split $\text{BP}\langle 2 \rangle \wedge Z_I \simeq B \vee \text{KV}$, where KV is the Eilenberg–MacLane spectrum for some graded vector space V , and H^*B has no free summands as a module over A ; then $h'_I: S \rightarrow B \vee \text{KV}$ is given by $* \vee f$ for some $f: S \rightarrow \text{KV}$. Since, for any spectrum X , $[\text{KV}, X] \cong \text{Hom}_A(A_*, H_* X)$

(the appropriate Adams spectral sequence collapses), we notice that the composite $S^j \circ h'_I$ is nonzero if and only if the induced map in homology is. Consider

$$\begin{array}{ccc} S^m & \xrightarrow{h'_I} B \vee KV & \xrightarrow{S^j \vee \text{Id}} \Sigma^{2(p-1)j} BP\langle 1 \rangle \wedge Z_I \\ & \downarrow \rho & \downarrow \rho \wedge \text{Id} \\ H\mathbb{Z}/p \wedge Z_I & \xrightarrow{\chi P^j \wedge \text{Id}} & \Sigma^{2(p-1)j} H\mathbb{Z}/p \wedge Z_I \end{array}$$

For dimensional reasons, since Z_I is a space, if $j > k$, $\chi P^i \circ \rho \circ h'_I \cong *$. Since reduction from $BP\langle 1 \rangle$ to $H\mathbb{Z}/p$ is a monomorphism in homology, we find that $S^i \circ h'_I$ is zero in homology. Hence $S^j \circ h'_I \cong *$ when $j > k$ so $d_{0*} h'_I = 0$.

3. Construction of $BP\langle 2 \rangle^k$

In this section we prove Theorem A for $n = 2$. The proof uses the fact that it has already been proved for $n = -1$. To get started we need the following lemma which encapsulates the role adaptive complexes play in the construction.

Lemma 3.1. *Let (T, h') be the adapted complex constructed in Section 2 and $h : DT \rightarrow K_0^2 = BP\langle 2 \rangle = X_0$ the dual of h' . Given parts (a), (b) and (c) of Theorem A for $q \leq t$, i.e. cofiber sequences*

$$K_{q+1}^2 \xrightarrow{i_{q+1}} X_{q+1} \xrightarrow{p_q} X_q \xrightarrow{e_q} \Sigma K_{q+1}^2$$

with $e_q i_q = d_q$, and given liftings $h_q : DT \rightarrow X_q$ for $q \leq t+1$. Then $\ker i_q^ \cap \ker h_q^* = 0$ for $q \leq t$.*

Proof. The proof is by induction on q . There are two base cases. If $q = 0$ the result follows since $i_0 = \text{Id}$. If $q = 1$, take $v \in \ker i_1^*$. Then $v = p_1^* w$ for some $w \in H^* X_0$. Now $\ker p_1^* = \text{Im } e_0^*$ implies that either $v = 0$ in which case we are done or else $w \notin \text{Im } e_0^* = \text{Im } d_0^* = \ker h_0^*$. The last equality is a consequence of Theorem 2.1. Thus $0 \neq h_0^* w = h_1 p_1^* w = h_1^* v$ where the second equality is a consequence of the fact that h_1 is a lifting of h_0 . Hence $\ker i_1^* \cap \ker h_1^* = 0$.

Now assume inductively that $\ker i_q^* \cap \ker h_q^* = 0$ for $q \leq s < t$. Take $v \in \ker i_{s+1}^*$. Then $v = p_s^* w$ for some $w \in H^* X_s$. Now $i_s^* w \in \ker d_{s-1}^* = \text{Im } d_s^*$ so choose $x \in H^* K_{s+1}^2$ such that $d_s^* x = i_s^* w$. Let $w' = w - e_s^* x$. Then $w' \in \ker i_s^*$ so by our induction hypothesis $h_s^* w' \neq 0$. But

$$h_s^* w' = h_{s+1}^* p_s^* (w - e_s^* x) = h_{s+1}^* v.$$

So $\ker i_{s+1}^* \cap \ker h_{s+1}^* = 0$. \square

Proof of Theorem A for $n = 2$. Let

$$\cdots \rightarrow Y_{q-1} \xrightarrow{p_{q-1}} Y_{q-1} \rightarrow \cdots \rightarrow Y_1 \rightarrow Y_0 = H\mathbb{Z}/p$$

be the tower whose homotopy inverse limit is $B(k)$, i.e. use $n = -1$ of Theorem A. We have the cofiber sequences

$$K_{q+1}^{-1} \xrightarrow{i_{q+1}} Y_{q+1} \xrightarrow{p_q} Y_q \xrightarrow{e_q} K_{q+1}^{-1}$$

and maps with the properties of Theorem A. One additional property of this case which we heavily depend on is given as Lemma 1.9 in [6], namely: for Z a finite complex, $m \leq 2p(k+1)-1$ and $\gamma: \Sigma^m DZ \rightarrow H\mathbb{Z}/p = Y_0$, any lifting of γ to Y_q lifts to Y_{q+1} . Let (Z, h') be the complex adapted to $M_2(k)$ constructed in Section 2. Set $T = \Sigma^{2pk+3} DZ$ and let $h: T \rightarrow BP\langle 2 \rangle$ be dual to h' .

We proceed by induction on q with the hypothesis:

$H(t)$: For $q \leq t+1$, there are spectra X_q and maps e_q, i_{q+1} for $q \leq t$ so that properties (a)-(d) of Theorem A hold. Additionally, there are maps $\theta'_q: X_q \rightarrow Y_q$ for $q \leq t+1$ so that for $q \leq t$

$$\begin{array}{ccccccc} K_{q+1}^2 & \xrightarrow{i_{q+1}} & X_{q+1} & \xrightarrow{p_q} & X_q & \xrightarrow{e_q} & \Sigma K_{q+1}^2 \\ \downarrow \theta_{q+1} & & \downarrow \theta'_{q+1} & & \downarrow \theta'_q & & \downarrow \theta_{q+1} \\ K_{q+1}^{-1} & \xrightarrow{i_{q+1}} & Y_{q+1} & \xrightarrow{p_q} & Y_q & \xrightarrow{e_q} & \Sigma K_{q+1}^{-1} \end{array} \quad (3.2)$$

is a commutative diagram of cofibration sequences. Also, h lifts to X_{t+1} .

Now X_0 is $BP\langle 2 \rangle$ and e_0 is $d_0: BP\langle 2 \rangle \rightarrow \bigvee_{j \geq k} \Sigma^{2(p-1)j} BP\langle 1 \rangle$. X_1 is defined to be the fiber of e_0 and setting i_0 to be the identity gives $e_0 i_0 = d_0$. $\theta'_0: BP\langle 2 \rangle \rightarrow H\mathbb{Z}/p$ is the usual reduction map and θ'_1 can be chosen to be any map satisfying (3.2), which completes the proof of $H(0)$.

$H(1)$ needs to be dealt with in a special manner. Define e_1 to fill in the following diagram of cofibrations, where the top square commutes because $d_1 \circ d_0 = 0$.

$$\begin{array}{ccc} \Sigma^{-1} BP\langle 2 \rangle = \Sigma^{-1} K_0^2 & \longrightarrow & * \\ \downarrow d_0 & & \downarrow \\ K_1^2 & \xrightarrow{d_1} & \Sigma K_2^2 \\ \downarrow i_1 & & \downarrow \\ X_1 & \xrightarrow{e_1} & \Sigma K_2^2 \end{array}$$

So by definition, $e_1 i_1 = d_1$. To get $e_1 \theta'_1 = \theta_2 e_1$ we may have to adjust e_1 a bit. Setting $\Delta = e_1 \theta'_1 - \theta_2 e_1$ we find that

$$\Delta i_1 = e_1 \theta'_1 - \theta_2 d_1 = d_1 \theta_1 - \theta_2 d_1 = 0.$$

Thus the top square in the following diagram commutes defining g .

$$\begin{array}{ccc}
K_1^2 & \longrightarrow & * \\
\downarrow i_1 & & \downarrow \\
X_1 & \xrightarrow{e_1\theta'_1 - \theta_2 e_1} & K_2^{-1} \\
\downarrow p_0 & & \parallel \\
BP\langle 2 \rangle & \dashrightarrow & K_2^{-1}
\end{array}$$

If $\rho: BP\langle 2 \rangle \rightarrow H$ is the reduction map, then, since $\rho^*: H^*H \rightarrow H^*BP\langle 2 \rangle$ is surjective and K_2^{-1} is a generalized Eilenberg-MacLane spectrum, g factors as

$$BP\langle 2 \rangle \xrightarrow{\rho = \theta_0} H = Y_0 \xrightarrow{g'} K_2^{-1}.$$

Define $e'_1 = e_1 - g'p_0: Y_1 \rightarrow K_2^{-1}$. Notice that in the tower defining $B(k)$ given in Theorem A, e_1 can be replaced by e'_1 since

$$e'_1 i_1 = (e_1 - g'p_0) i_1 = e_1 i_1 = d_1.$$

This replacement may require adjustment of the higher e_i 's but this can be done as in the original proof. So

$$e'_1 \theta'_1 - \theta_2 e_1 = \Delta - g'p_0 \theta_1 = \Delta - g' \theta'_0 p_0 = \Delta - \Delta = 0.$$

To get the lifting of h_1 notice that $\theta'_1 h_1$ lifts, so $\theta_2 e_1 h_1 = e_1 \theta'_1 h_1 = 0$. T was chosen so that $BP\langle 0 \rangle^* T \rightarrow H\mathbb{Z}/p^* T$ is a monomorphism, hence $e_1 h_1 = 0$ and h_1 lifts to $h_2: T \rightarrow X_2$, completing the verification of $H(2)$.

Now assume $H(t-1)$ with $t \geq 2$. This proof of the induction step is essentially that of [6]. Since $t+1 \geq 3$, there is a map $s_{t+1}: K_{t+1}^{-1} \rightarrow K_{t+1}^2$ such that $s_{t+1} \theta_{t+1}$ is the identity. Set

$$e_t = s_{t+1} e_t \theta'_t: X_t \rightarrow \Sigma K_{t+1}^2$$

and let X_{t+1} be the fiber of e_t . Then by our induction hypothesis,

$$e_t i_t = s_{t+1} e_t \theta'_t i_t = s_{t+1} e_t i_t \theta_t = s_{t+1} d_t \theta_t = s_{t+1} \theta_{t+1} d_t = d_t.$$

To lift h_t to X_{t+1} , notice that $\theta'_t h_t$ is a lifting of $\theta'_0 h$. By Lemma 1.7 of [6] this must lift even further so we must have $e_t \theta'_t h_t = 0$ so $0 = s_{t+1} e_t \theta'_t h_t = s_{t+1} \theta_{t+1} e_t h_t = e_t h_t$ and h_t lifts. To get the diagram (3.2) we need to show

$$\begin{array}{ccc}
X_t & \xrightarrow{e_t} & \Sigma K_{t+1}^2 \\
\downarrow \theta'_t & & \downarrow \theta_{t+1} \\
Y_t & \longrightarrow & \Sigma K_{t+1}^{-1}
\end{array}$$

commutes and then simply define θ'_{t+1} to satisfy (3.2). Since K_{t+1}^{-1} is an Eilenberg-MacLane spectrum, it suffices to show $\theta_{t+1} e_t = e_t \theta'_t$ in cohomology. Lemma 3.1

indicates that it is sufficient to show $i_t^* e_t^* \theta_{t+1}^* = i_t^* \theta_t^* e_t^*$ and $h_t^* e_t^* \theta_{t+1}^* = h_t^* \theta_t^* e_t^*$. However

$$\theta_{t+1} e_t i_t - e_t \theta_t' i_t = \theta_{t+1} d_t - e_t i_t \theta_t = \theta_{t+1} d_t - d_t \theta_t = 0$$

and an earlier argument showed $e_t h_t = 0 = e_t \theta_t' h_t$. Finally, property (d) of Theorem A holds here because

$$(\theta_{t+1})_*: (K_{t+1}^2)_* Z \rightarrow (K_{t+1})_* Z$$

is injective. This completes the proof of the inductive step and the theorem. \square

To finish this section, we need to complete the proof of Theorem B begun in the introduction. Recall that $BP\langle 2 \rangle^k$ is defined to be $\text{holim}_q X_q$. We have already shown that if Z is a spectrum, $w: BP\langle 2 \rangle^k \rightarrow BP\langle 2 \rangle$ the map from $\text{holim}_q X_q$ to X_0 and $m \leq 2p(k+1)-1$ then $\{\text{Im } w_*: BP\langle 2 \rangle_m^k Z \rightarrow BP\langle 2 \rangle_m Z\} \subseteq \bigcap_{j>k} \ker S_*^j$ and we need to show that if $H\mathbb{Z}_p^{\wedge} Z$ is an F_p vector space then $\text{Im } w_* = \bigcap_{j>k} \ker S_*^j$.

Proof. Smash all of the diagrams used to construct $BP\langle 2 \rangle^k$ with Z . Take any $f: S^m \rightarrow BP\langle 2 \rangle \wedge Z$ with the property that $f \in \bigcap_{j>k} \ker S_*^j \subseteq BP\langle 2 \rangle_m Z$ and $m \leq 2p(k+1)-1$. The first condition on f guarantees that f lifts to $f_1: S^m \rightarrow X_1 \wedge Z$. Let \bar{f}_1 be the composite

$$S^m \xrightarrow{f_1} X_1 \wedge Z \xrightarrow{\theta_1' \wedge \text{Id}} Y_1 \wedge Z.$$

Then by Lemma 1.9 of [6], \bar{f}_1 lifts to $\bar{f}_2: S^m \rightarrow Y_2 \wedge Z$ so that $(e_1 \wedge \text{Id}_Z) \circ \bar{f}_1 = 0$. Writing Z to denote the identity map on Z , the commutativity of (3.2) tells us that

$$0 = (e_1 \wedge Z) \circ (\theta_1' \wedge Z) \circ f_1 = (\theta_2 \wedge Z) \circ (e_1 \wedge Z) \circ f_1.$$

Since $H\mathbb{Z}_p^{\wedge} Z$ is an F_p vector space, this means that $(e_1 \wedge Z) \circ f_1 = 0$ and so f_1 lifts to f_2 . A similar argument shows that f_2 and higher lifts can always be lifted. These arguments don't need $H\mathbb{Z}_p^{\wedge} Z$ an F_p vector space, since K_s^2 is a wedge of $H\mathbb{Z}/p$'s for $s \geq 3$, but do use the fact that K_s^2 is a wedge summand of K_s^{-1} for $s \geq 3$. Thus f lifts all the way up the tower to give $f': S^m \rightarrow BP\langle 2 \rangle^k \wedge Z$ with $w \circ f' = f$.

4. Spacelike spectra and maps to Brown–Gitler spectra

Proof of Corollary D. Throughout this proof, all pairs (s, t) should be assumed to satisfy $s - t \geq n - 2p(k+1) + 2$. The first thing to notice is that since Y is a finite spectrum, $[Y, BP\langle 2 \rangle^k] \cong [s, DY \wedge BP\langle 2 \rangle^k]$, and we can replace $E_r^{**}(Y, BP\langle 2 \rangle^k)$ by $E_r^{**}(DY \wedge BP\langle 2 \rangle^k)$. Let $f: \Sigma^n DZ \rightarrow Y$ be the map making Y spacelike. Then $Df: DY \rightarrow \Sigma^{-n} Z$ has the property that $Df_*: H_* DY \rightarrow H_* \Sigma^{-n} Z$ is a monomorphism. Df induces a map $i: E_1^{s,t}(DY \wedge BP\langle 2 \rangle^k) \rightarrow E_1^{s,t}(\Sigma^{-n} Z \wedge BP\langle 2 \rangle^k)$ which is monic for $s \geq 3$ since Df_* is monic. Z being a space, Corollary C tells us that all differentials in $E_r^{**}(\Sigma^{-n} Z \wedge BP\langle 2 \rangle^k)$ are zero except possibly for d_1 and d_2 .

The proof is by induction on r in the assertion that for $s - t \geq n - 2(k + 1) + 2$,

$$E_3^{s,t}(DY \wedge BP\langle 2 \rangle^k) \cong E_r^{s,t}(DY \wedge BP\langle 2 \rangle^k), \text{ for all } r \geq 3 \quad (\text{A})$$

and

$$i: E_r^{s,t}(DY \wedge BP\langle 2 \rangle^k) \rightarrow E_r^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k) \text{ is monic for } s \geq 3. \quad (\text{B})$$

The base case is $r=3$ where (A) is immediate. As noted above, $i: E_1^{s,t}(DY \wedge BP\langle 2 \rangle^k) \rightarrow E_1^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k)$ is monic for $s \geq 3$. Now consider the first differential. In $E_1^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle)$, d_1 is zero on $E_1^{s,t}$ for $s \geq 2$ because in that range K_{s+1}^2 is a wedge summand of K_{s+1}^{-1} , and maps to the original Brown-Gitler tower always lift. Said more explicitly, $x \in E_1^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k)$ is a map $x: S^{s-t-n} \rightarrow \Sigma^{-n}Z \wedge K_s^2$.

$$\theta_{s+1}d_1x = \theta_{s+1}e_{s,t}i_sx = e_{s+1}\theta'_{s+1}i_sx = 0$$

because $\theta'_{s+1}i_sx = S^{s-t-n} \rightarrow \Sigma^{-n}Z \wedge Y_{s+1}$ lifts. However $s+1 \geq 3$ implies that θ_{s+1} is an inclusion of a wedge summand so $d_1x = 0$. Thus

$$E_1^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k) \cong E_2^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k).$$

Therefore d_1 is zero on $E_1^{s,t}(DY \wedge BP\langle 2 \rangle^k)$ for $s \geq 3$ since $0 = d_1ix = \text{id}_1x$ and i is monic. A similar argument shows d_2 is zero on $E_2^{s,t}$ for $s \geq 1$ so

$$E_3^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k) \cong E_2^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k) \text{ for } s \geq 3 \text{ and}$$

$$E_3^{s,t}(DY \wedge BP\langle 2 \rangle^k) \cong E_2^{s,t}(DY \wedge BP\langle 2 \rangle^k) \text{ for } s \geq 5 \text{ and}$$

$$i: E_3^{s,t}(DY \wedge BP\langle 2 \rangle^k) \rightarrow E_3^{s,t}(\Sigma^{-n}Z \wedge BP\langle 2 \rangle^k) \text{ is monic for } s \geq 3.$$

For the induction step, assume both are true for some $n \geq 3$. Since d_nx has filtration ≥ 3 for any $x \in E_n^{s,t}(DY \wedge BP\langle 2 \rangle^k)$, $0 = d_nix = \text{id}_n x$. Since i is monic in this range, $d_nx = 0$. Thus both $E_{n+1}^{s,t}$ terms are isomorphic to their respective $E_n^{s,t}$ terms and i remains monic on $E_{n+1}^{s,t}$ for $s \geq 3$. This completes the induction step. \square

One final comment is in order. [6] used these spectral sequences to construct pairings $BP\langle 1 \rangle^i \wedge BP\langle 1 \rangle^j \rightarrow BP\langle 1 \rangle^{i+j}$ and to prove the uniqueness of $BP\langle 1 \rangle^k$. I have not been able to prove the analogous results for the $BP\langle 2 \rangle^k$ case because of the extra differential involved.

Acknowledgement

This paper stems from my Ph.D. thesis done under the supervision of John Jones at the University of Warwick. I want to thank him for suggesting the project to me and also the Natural Sciences and Engineering Research Council of Canada and the University of Western Ontario for providing me with support and a place to work while writing this paper. I also want to thank Paul Goerss for warning me of certain pitfalls and Mark Mahowald for helpful comments particularly regarding style.

References

- [1] E.H. Brown and S. Gitler, A spectrum whose cohomology is a certain cyclic module over the Steenrod algebra, *Topology* 12 (1973) 283–295.
- [2] E.H. Brown and F.P. Peterson, On the stable decomposition of $\Omega^2 S^{r+2}$, *Trans. Amer. Math. Soc.* 243 (1978) 287–298.
- [3] R.L. Cohen, Odd primary infinite families in stable homotopy theory, *Memoirs Amer. Math. Soc.* 242 (1981).
- [4] P.G. Goerss, Results on Brown-Gitler spectra, Thesis, M.I.T., 1983.
- [5] P.G. Goerss, in: *Proc. 1985 Arcata Confer. Brown-Gitler spectra and applications*, to appear.
- [6] P.G. Goerss, J.D.S. Jones and M.E. Mahowald, Some generalized Brown-Gitler spectra, *Trans. Amer. Math. Soc.* 294 (1986) 113–132.
- [7] R. Kane, Operations in connective K -theory, *Memoirs Amer. Math. Soc.* 254 (1981).
- [8] J. Klippenstein, Applications of the universal coefficient theorem for connective K -theory, Thesis, University of Warwick, 1985.
- [9] M.E. Mahowald, bo resolutions, *Pac. J. Math.* 92 (1981) 365–383.
- [10] D. Shimamoto, An integral version of the Brown-Gitler spectrum, *Trans. Amer. Math. Soc.* 283 (1984) 383–421.